




Epermenia alba *punctella* No common name



High counts of:

- 1 - Madison - 2019-05-05
- 1 - Madison - 2019-05-19
- 1 - Madison - 2020-03-13

Status Rank
NC US NC Global



FAMILY: Epermeniidae SUBFAMILY: [Epermeniinae] TRIBE: [Epermeniini]
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: In the Northeast, larvae and pupae have been found beginning in early July, with adults emerging from early July to early August (Eiseman 2019).

HABITAT:

FOOD: This species has been reported to use several taxa in the Apiaceae (*Daucus*, *Heracleum*, *Ligusticum*, *Osmorhiza*), as well as *Aralia* in the Araliaceae (Eiseman 2019, 2019a). Prentice (1966) also reported it to use conifers in Canada. The reported hosts include Queen-Anne's-Lace (*Daucus carota*), Sweet-cicely (*Osmorhiza claytonii*), Norway Spruce (*Picea abies*), White Spruce (*P. glauca*) and Balsam Fir (*Abies balsamea*).

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS:

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: