




# *Diasemiopsis leodocusalis* Brindled Shawl Moth



FAMILY: Crambidae SUBFAMILY: Pyraustinae TRIBE: Spilomelini  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS:

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: This species has complex but distinctive wing patterning. The forewing has a gray ground color that is overlain with a prominent white postmedial band that runs perpendicular from the costa and ends about one-third of the way inward. A second, whitish, irregular band originates on the inner margin just beyond the middle and runs obliquely outward towards the end of the postmedial band. Several blackish streaks or blotches are present on the wing. These include a diffuse blotch immediately apical to the postmedial band, a longitudinal streak immediately basal to the postmedial band, a short, diffuse streak immediately apical to the median white band, and a longer streak immediately basal to the median band. A white subcostal spot is sometimes evident in the middle of the wing. The entire wing is cross-marked with numerous short tan striations, and the costa has a series of tan dots from the postmedial band to the wing base. The fringe is checkered with a series of white and dark brown marks.

The hindwing is concave near the apex and arcuate. It is boldly marked with a series of irregular and alternating black and white bands, beginning with a narrow black band at the base that is followed by larger bands of alternating white and black color. The final band adjoins a sinuous black line that is bordered outwardly by a thin white line. The terminal third of the wing has a blackish shade and the fringe is checkered with alternating white and dark brown marks. The blackish portions of the wing are cross-marked with numerous short tan striations as seen in the forewing.

DISTRIBUTION: *Diasemiopsis leodocusalis* occurs in Central and South America, the West Indies and throughout much of the southeastern U.S., where the range extends from Long Island, New York southward to southern Florida and westward to eastern Texas, Arkansas and western Tennessee. As of 2023, all of our records are from the Piedmont and southern Coastal Plain.

FLIGHT COMMENT: The adults fly year-round in Florida and from July through December elsewhere. As of 2023, our records extend from early-August through mid-November.

HABITAT: Specimens have been collected from xeric, mesic and hydric habitats, including a Long-leaf Pine forest and a brackish marsh.

FOOD: The hosts are undocumented.

OBSERVATION\_METHODS: The adults are attracted to lights.

NATURAL HERITAGE PROGRAM RANKS: GNR [S2-S3]

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: