




Condylolomia participialis Drab Condylolomia Moth



FAMILY: Pyralidae SUBFAMILY: Chrysauginae TRIBE: [Chrysaugini]
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: We have records from a wide variety of habitats, including Northern Hardwood and Spruce-Fir Forests, high elevation grasslands, mountain bogs and alluvial forests, and residential areas in the Piedmont

FOOD: Larval host plants include members of the Myricaceae (Covell, 1984). Myrica gale is specifically mentioned by BugGuide (accessed 2020-10-29). In North Carolina, however, Myrica gale is extremely rare and bayberries and wax myrtles occur mainly in the Coastal Plain where this moth appears to be scarce or absent.

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: GNR SU

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: