




# *Lobocleta plemyraria* Straight-lined Wave Moth



FAMILY: Geometridae SUBFAMILY: Sterrhinae TRIBE: Sterrhini  
TAXONOMIC\_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Almost all of our records for this species come from dry-xeric sand ridges in the Coastal Plain. One Piedmont record comes from dry woodlands on a monadnock.

FOOD: Reported to feed on *Euphorbia albomarginata*, a Southwestern species (Robinson et al., 2010) and *Chamaesyce*, another western spurge (Powell & Opler, 2009). Xerophytic members of the Euphorbiaceae are also common members of the sandhill communities of the Southeast and may be used here as well.

OBSERVATION\_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G2G4 S2S3

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: