




Renia sobrialis Sober Renia Moth



FAMILY: Erebidae SUBFAMILY: Herminiinae TRIBE:
TAXONOMIC_COMMENTS:

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS:

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS: The ground color is a dark, smoky gray with little or no contrasts (Forbes, 1954). The antemedian and postmedian are inconspicuous but darker than the ground color and either dentate or irregular. In our specimens, the subterminal often consists of narrow white dots. The orbicular and reniform are yellowish or orange; the reniform can have large, dark central spots that sometimes forms a connected lunule.

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: The majority of our records come from Longleaf Pine savannas, seeps, bog, shorelines, and other open, wet, herb-dominated habitats

FOOD: Probably feeds on dead leaves and other organic debris like other members of this genus

OBSERVATION_METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S4S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: