
Acronicta fragilis Fragile Dagger



FAMILY: Noctuidae SUBFAMILY: Acronictinae TRIBE:

TAXONOMIC_COMMENTS: One of 74 species in this genus found in North America north of Mexico (Schmidt and Anweiler, 2020), 42 of which have been recorded in North Carolina. <i>Acronicta fragilis</i> is a member of Species Group II of Forbes (1954), which includes 17 additional species in North Carolina.

FIELD GUIDE DESCRIPTIONS: Covell (1984); Beadle and Leckie (2012)

ONLINE PHOTOS:

TECHNICAL DESCRIPTION, ADULTS: Forbes (1954); Schmidt and Anweiler (2020)

TECHNICAL DESCRIPTION, IMMATURE STAGES:

ID COMMENTS:

DISTRIBUTION: Please refer to the dot map.

FLIGHT COMMENT: Please refer to the flight charts.

HABITAT: Schmidt and Anweiler (2020) describe this species as associated with cool, temperate, moist-to-wet woodland habitats. Our records come from alluvial and cove forests at low elevations in the Mountains and more commonly in high elevation forests.

FOOD: Larvae feed on a wide range of hardwood trees and shrubs, including Alder, Apple, Beech, Birch, Cherry, Mountain Ash, Plum, Rose, Serviceberry, and Willow (Wagner et al., 2011). Species in the Rosaceae seem to be preferred, with members of the Betulaceae also commonly used (Schmidt and Anweiler, 2020).

OBSERVATION METHODS:

NATURAL HERITAGE PROGRAM RANKS: G5 S4S5

STATE PROTECTION: Has no legal protection, although permits are required to collect it on state parks and other public lands.

COMMENTS: